2012

Current Exercise Behaviors of Breast Cancer Patients Diagnosed with Chemotherapy-Induced Peripheral Neuropathy

Karen Y. Wonders
Wright State University

Daniel G. Drury
Gettysburg College

Follow this and additional works at: <https://cupola.gettysburg.edu/healthfac>

Part of the [Oncology Commons](#)

Share feedback about the accessibility of this item.

Wonders KY, Drury DG (2012) Current Exercise Behaviors of Breast Cancer Patients Diagnosed with Chemotherapy-induced Peripheral Neuropathy. *Journal of Integrative Oncology* 1:103. <http://dx.doi.org/10.4172/jio.1000103>

This is the publisher's version of the work. This publication appears in Gettysburg College's institutional repository by permission of the copyright owner for personal use, not for redistribution. Cupola permanent link: <https://cupola.gettysburg.edu/healthfac/17>

This open access article is brought to you by The Cupola: Scholarship at Gettysburg College. It has been accepted for inclusion by an authorized administrator of The Cupola. For more information, please contact cupola@gettysburg.edu.

Current Exercise Behaviors of Breast Cancer Patients Diagnosed with Chemotherapy-Induced Peripheral Neuropathy

Abstract

Introduction: Chemotherapy-induced peripheral neuropathy (CIPN) is a common, dose-limiting effect of cancer therapy. The neuropathic pain associated with CIPN often has negative implications on an individual's quality of life (QOL) and has long been recognized as one of the more difficult types of pain to treat. Treatment of neuropathic pain due to CIPN often requires a multidisciplinary approach, with much attention focused on the use of pharmacological therapies. However, in most instances, these agents have been shown to have additional negative side effects for cancer patients. Thus, other interventions that address the symptoms of CIPN should be considered. One such possible intervention is exercise rehabilitation, which has previously been reported effective in attenuating numerous cancer treatment-related toxicities and enhancing the QOL of patients. However, to our knowledge, there have been no published clinical trials examining the role of exercise in preserving neurological function following chemotherapy. As such, the purpose of this investigation was to examine the current exercise habits of breast cancer patients who are diagnosed with CIPN and the impact on pain and QOL.

Methods: 300 women listed in the Breast Cancer Registry of Greater Cincinnati database were recruited by mail and asked to complete three questionnaires (McGill QOL, Leeds Assessment of Neuropathic Symptoms and Signs, and Current Exercise Behaviors). Data was analyzed at the 0.05 level of significance using a student's t-test and a Pearson's product moment correlation.

Results: 134 completed surveys were returned and analyzed (44.6% response rate). Overall, QOL and exercise behaviors were moderately correlated ($r = 0.56$). Patients reported exercising an average of 2.3 d/wk and an overall QOL of 4.7. Of the patients completing the recommended amount of physical activity (EX, $n = 21$), QOL was 6.3, which was significantly higher than patients who did not meet these recommendations (SED, $n = 113$, $p < 0.001$). Likewise, only 15% of EX patients reported experiencing pain compared to 72% of SED patients ($p < 0.001$). **Conclusions:** Based on these data, it seems likely that an exercise intervention would be successful in attenuating symptoms of CIPN and improving the overall QOL of breast cancer patients.

Keywords

Chemotherapy-induced peripheral neuropathy, Exercise behaviors, Quality of life, Pain

Disciplines

Medicine and Health Sciences | Oncology

Research Article

Current Exercise Behaviors of Breast Cancer Patients Diagnosed with Chemotherapy-induced Peripheral Neuropathy

Karen Y. Wonders^{1,2*} and Daniel G. Drury³¹Department of Health, Physical Education and Recreation, Wright State University, 316 Nutter Center, 3640 Colonel Glenn Hwy, Dayton, OH 45435, USA²Maple Tree Cancer Alliance, 106 Peach Orchard Ave, Dayton, OH, 45419, USA³Gettysburg College, Department of Health Sciences, 300 N. Washington Street, Gettysburg, PA 17325, USA

Abstract

Introduction: Chemotherapy-induced peripheral neuropathy (CIPN) is a common, dose-limiting effect of cancer therapy. The neuropathic pain associated with CIPN often has negative implications on an individual's quality of life (QOL) and has long been recognized as one of the more difficult types of pain to treat. Treatment of neuropathic pain due to CIPN often requires a multidisciplinary approach, with much attention focused on the use of pharmacological therapies. However, in most instances, these agents have been shown to have additional negative side effects for cancer patients. Thus, other interventions that address the symptoms of CIPN should be considered. One such possible intervention is exercise rehabilitation, which has previously been reported effective in attenuating numerous cancer treatment-related toxicities and enhancing the QOL of patients. However, to our knowledge, there have been no published clinical trials examining the role of exercise in preserving neurological function following chemotherapy. As such, the purpose of this investigation was to examine the current exercise habits of breast cancer patients who are diagnosed with CIPN and the impact on pain and QOL.

Methods: 300 women listed in the Breast Cancer Registry of Greater Cincinnati database were recruited by mail and asked to complete three questionnaires (McGill QOL, Leeds Assessment of Neuropathic Symptoms and Signs, and Current Exercise Behaviors). Data was analyzed at the 0.05 level of significance using a student's t-test and a Pearson's product moment correlation.

Results: 134 completed surveys were returned and analyzed (44.6% response rate). Overall, QOL and exercise behaviors were moderately correlated ($r = 0.56$). Patients reported exercising an average of 2.3 d/wk and an overall QOL of 4.7. Of the patients completing the recommended amount of physical activity (EX, $n = 21$), QOL was 6.3, which was significantly higher than patients who did not meet these recommendations (SED, $n = 113$, $p < 0.001$). Likewise, only 15% of EX patients reported experiencing pain compared to 72% of SED patients ($p < 0.001$). Conclusions: Based on these data, it seems likely that an exercise intervention would be successful in attenuating symptoms of CIPN and improving the overall QOL of breast cancer patients.

Keywords: Chemotherapy-induced peripheral neuropathy; Exercise behaviors; Quality of life; Pain

Introduction

Chemotherapy-induced peripheral neuropathy (CIPN) is a common ailment among those taking chemotherapeutic drugs. This condition is characterized by damage to the nervous system that is a direct result of the medications associated with chemotherapy. Little is known about the mechanisms responsible for the development of CIPN [1]. The peripheral toxicity involved with CIPN is specific to each chemotherapy drug class, and in most cases, appears to be dose and duration dependent, however it can evolve even after a single drug application [2]. The chemotherapy agents Docetaxel, Paclitaxel, or Vinorelbine are used to treat breast cancer, and often result in damage to central nervous system pain pathways, resulting in neuropathic pain [3-6]. The presence and severity of neuropathic pain is often shown to be associated with impairments in walking, general activities, sleep, work, mood, enjoyment of life, and relationships with others [7,8].

Much effort has been made to explore pharmacological therapies to reduce CIPN. Some of these therapies provide modest improvements in neurological function. However, in most instances, these agents have been shown to have additional negative side effects for cancer patients, such as cardiac conduction defects, abnormalities in antidiuretic hormone secretion, loss of balance, swelling in extremities, tremor, and increased chemotherapy resistance [9-11]. Thus, other interventions that address the symptoms of CIPN should be considered. One intervention that has the potential of preventing or alleviating CIPN is exercise reha-

bilitation. Exercise has been shown to have a local effect on peripheral nerves, inducing changes in both the vasculature and metabolic systems [12]. Short-term exercise stimulates endothelium-dependent vasodilatation and endoneurial blood flow [13]. Long-term exercise has a positive effect on oxygen delivery, as increased blood flow exposes blood vessels to shear stress, which augments vasodilatation [13]. Several studies have illustrated the beneficial effects of exercise in attenuating numerous cancer treatment-related toxicities and enhancing the quality of the lives of patients [14-21]. In addition, numerous studies on the effect of exercise in populations with diabetic peripheral neuropathy have produced promising results [22-24]. However, published clinical trials examining the role of exercise in preserving neurological function following chemotherapy are limited. Since breast cancer survivors represent the largest population of cancer survivors in the United States,

***Corresponding author:** Karen Y. Wonders, Wright State University, Department of Health, Physical Education and Recreation, Wright State University, 316 Nutter Center, 3640 Colonel Glenn Hwy, Dayton, OH 45435, USA, Tel: 937-775-2637; Fax: 937-775-4252; E-mail: karen.wonders@wright.edu

Received March 08, 2012; Accepted April 25, 2012; Published April 27, 2012

Citation: Wonders KY, Drury DG (2012) Current Exercise Behaviors of Breast Cancer Patients Diagnosed with Chemotherapy-induced Peripheral Neuropathy. J Integr Oncol 1:103. doi:10.4172/jio.1000103

Copyright: © 2012 Wonders KY, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

the purpose of this investigation was to examine the current exercise behaviors of breast cancer patients who were diagnosed with CIPN and to determine the resulting impact on their pain and quality of life.

Methods

Subjects

This investigation used a non-probability sample of convenience. Women who met the eligibility criteria and were listed in the Breast Cancer Registry of Greater Cincinnati (BCRGC) database were recruited by mail in the fall of 2010. The eligibility criteria included women who (a) have recently received a clinical diagnosis of CIPN, and (b) were currently taking the chemotherapy agents Docetaxel, Paclitaxel, or Vinorelbine, and (c) were able to read and write English. The BCRGC is a database established by the University of Cincinnati, Department of Environmental Health for the purpose of supporting studies that focus on breast cancer issues. Women who were breast cancer survivors and were located in the Greater Cincinnati area may become members by signing a consent form and completing a registration form that includes questions specific to the detection, diagnosis, and treatment of their disease. A total of 300 women listed in the BCRGC met the eligibility criteria of the study. All procedures were approved by the Wright State University Institutional Review Board prior to data collection.

Data collection

The 300 women who met the eligibility criteria and agreed to be contacted received a packet that included a cover letter, the McGill Quality of Life questionnaire [25], the Leeds Assessment of Neuropathic Symptoms and Signs questionnaire (LANSS) [26], a questionnaire to measure current exercise behaviors (Reigle), and a self-addressed, stamped envelope. Women were asked to return the completed questionnaires within two weeks of receiving the packet. Women who did not respond within two weeks were sent a reminder postcard.

Data analysis

Descriptive statistics have been computed as means and standard deviations. A Pearson's product moment correlation was used to determine the relationship between QOL and current exercise behaviors for the group as a whole. A student's T-test was used to compare the overall QOL ratings, as well as the prevalence of pain ratings, among patients who received the recommended amount of exercise compared to those who did not. All data was analyzed at the 0.05 level of significance.

Results

Of the 300 surveys sent out, 134 completed surveys were returned and analyzed, giving a 44.6% response rate. The characteristics of the patients are summarized in Table 1. The distribution of QOL was normal. On average, patients reported exercising 2.3 d/wk (approximately 69 min/wk) and rated their overall quality of life (QOL) as 4.7 on a 10-point scale. CIPN impacted routine activities, functions, and behaviors in the areas of domestic, work, and social/leisure life. A moderate correlation was measured between QOL and current exercise behaviors for the group as a whole ($r = 0.56$). Table 2 summarizes the results of the Exercise Behaviors Survey for all respondents. Table 3 presents the results of the LANSS survey for all respondents.

Responses were analyzed to determine the percentage of patients completing the recommended amount of physical activity (at least 150 min moderate-intense ex/wk and muscle strengthening activity 2 d/wk

Age at time of survey (yrs)	Age at time of diagnosis (yrs)	Education Level	Race	Length of Cancer Treatment
54.9 ± 2.4	50.4 ± 5.7	36 = high school 68 = college 30 = postgrad	White = 105 Black = 21 Hispanic = 6 Other = 2	4.5 ± 6.4 mo's

Table 1: Subject Characteristics. Values are means ± SE.

Type of Exercise	Frequency/wk
Increase flexibility of arms and shoulders	2.39 ± 0.10
Increase strength of arms and shoulders	2.16 ± 0.10
Increase flexibility of legs	2.48 ± 0.11
Increase strength of legs	2.45 ± 0.10
Increase strength and flexibility of upper body	2.16 ± 0.10
Increase strength and flexibility of lower body	2.35 ± 0.10
Balance exercises	2.03 ± 0.11
Resistance training	1.97 ± 0.09

Table 2: Results from Exercise Behaviors Survey. Values are means ± SE.

Questions	% of respondents answering Yes
Would you describe your pain as pricking, tingling, pins and needles?	51 ± 1.2%
Does the skin in the painful areas look different than normal?	17.9 ± 0.5%
Is the skin sensitive to touch?	33.3 ± 1.7%
Does your pain come on suddenly in bursts when you are still?	42.3 ± 1.1%
Do you feel that skin temperature in the painful area has changed?	20.5 ± 2.1%
Does stroking the affected area with cotton wool produce pain?	9.2 ± 0.3%

Table 3: Results from LANSS Survey. Values are means ± SE.

[27]. Approximately 15.6% of the patients surveyed reported meeting these recommended levels of physical activity (EX, $n = 21$). This group of patients reported an overall QOL of 6.3. As demonstrated in (Figure 1), this was significantly higher than those patients who did not meet these recommendations (SED, $n = 113$), who reported an average overall QOL of 3.4 ($p < 0.001$). Similarly, only 15% of EX patients reported experiencing pain compared to 72% of SED patients ($p < 0.001$) (Figure 2). The pain was frequently described as "tingling" and "numbness" and localized to "hands and feet". Several respondents ($n = 46$) indicated that this pain was "better when I exercise".

Discussion

The major finding in this investigation was that breast cancer patients with CIPN who received the recommended levels of physical activity reported a significantly higher overall QOL and experienced significantly less pain than their sedentary counterparts. CIPN affected breast cancer patients' routine activities, functions, and behaviors. While it is unknown if the sedentary individuals abstained from exercise because of their pain, the respondents who stated that their pain was "better when I exercise" indicates that it seems likely that an exercise intervention would be successful in attenuating symptoms of CIPN and improving the overall QOL of breast cancer patients.

With neuropathy, muscle mass atrophies and leads to significant decreases in muscular strength [28]. This decline in strength appears to be slow and progressive. It also appears to affect distal muscle groups more so than proximal muscles. Researchers have indicated that this muscle weakness translates into impaired motor performance skills and a reduced exercise capacity [29]. However, several studies have reported improvements in muscular strength following moderate resistance exercise programs in patients with hereditary motor and sensory neuropathies [30,31], as well as diabetic neuropathies, and those associated with fibromyalgia and chronic fatigue [32-37]. In light of these findings, many researchers recommend that exercise training serve as an important component in the comprehensive treatment plan for patients with peripheral neuropathy [33,34,38]. Moderate-to-intense strength training and aerobic exercise appears to be well tolerated by these patients [38], and is associated with improvements in motor function and nerve conduction velocity [32,35], as well as improved muscle reinnervation and increased axon regeneration [39]. In addition, one investigation reported that low intensity treadmill exercise promoted Schwann cell proliferation in the injured peripheral nerve [40]. In light of these findings, it is feasible to assume that an individual who has experienced a reduction in muscular strength and functional ability due to CIPN may experience similar improvements following an exercise program.

It is not uncommon for individuals with cancer to experience pain. This pain may be triggered by the disease itself, or by its associated treatments. The direct effect of the tumor accounts for about 70% of the pain related to cancer, while approximately 15% of cancer pain is related to its treatments [41]. The pain associated with CIPN has long been recognized as one of the more difficult types of pain to treat. However, as explained earlier, it is possible that exercise may have a beneficial effect on pain. As illustrated by the respondents in the cur-

rent study, those individuals who exercised experienced significantly less pain than their sedentary counterparts. Likewise, it was noted that pain became "better" upon exercise. This finding is supported by several studies conducted on healthy populations, whereby pain thresholds and tolerance levels have been reported to increase both during and following exercise. In addition, intensity ratings of pain appear to decrease following exercise [42,43].

While literature examining the role of exercise on symptoms of pain related to CIPN is lacking, we believe that the existing tangential research supports the use of exercise as a means to attenuate general cancer related pain. Exercise during cancer treatment has been shown to have a beneficial effect on pain for breast cancer patients [44-47]. In a similar study to ours, Liou et al. [45] surveyed the daily physical activity levels of individuals who were undergoing chemotherapy, but were not diagnosed with CIPN. These authors reported that those patients who received the recommended 150 min/wk of moderate activity reported significantly less symptoms of pain than their sedentary counterparts.

In light of the findings and the literature surrounding this topic, it seems feasible to assume that a supervised exercise program would provide a beneficial effect on quality of life and pain associated with CIPN in breast cancer patients. More follow up investigations are warranted.

Limitations

The most apparent limitation in the present study is the low response rate from the breast cancer patients (44.6%). However, in survey research, no single response rate is considered a standard [48]. Mail surveys typically have lower response rates than other types of surveys [49]. In the present study, non respondents were sent a reminder postcard in an attempt to increase the response rate; however, additional measures could have also been employed, such as personal contact. In addition, due to the lack of existing literature examining the effects of exercise on symptoms of CIPN, the authors had to look to the effects of exercise on symptoms of peripheral neuropathy in other chronic diseases (i.e., diabetes) to make its generalizations. Thus, it is possible that an exercise program may not produce similar results in a breast cancer patient. Likewise, since the location of cancer will vary the symptoms related to CIPN, it is possible that the exercise response would produce a different outcome in another form of cancer.

References

- Kaley TJ, Deangelis LM (2009) Therapy of chemotherapy-induced peripheral neuropathy. *Br J Haematol* 145: 3-14.
- Quasthoff S, Hartung HP (2002) Chemotherapy-induced peripheral neuropathy. *J Neurol* 249: 9-17.
- Horowitz SH (2007) The diagnostic workup of patients with neuropathic pain. *Med Clin North Am* 91: 21-30.
- National Health & Medical Research Council (1999) Acute pain management: Scientific evidence Australia. Commonwealth of Australia.
- Verstappen CCP, Heimans JJ, Hoekman K, Postma TJ (2003) Neurotoxic complications of chemotherapy in patients with cancer: Clinical signs and optimal management. *Drugs* 63: 1549-1563.
- Visovsky C (2003) Chemotherapy-induced peripheral neuropathy Nursing Perspectives. *Cancer Investigation* 21: 439-451.
- Tolle T, Xu X, Sadosky AB (2006) Painful diabetic neuropathy: a cross-sectional survey of health state impairment and treatment patterns. *J Diabetes Complications* 20: 26-33.
- Zelman DC, Gore M, Dukes E, Tai KS, Brandenburg N (2005) Validation of a modified version of the brief pain inventory for painful diabetic peripheral neuropathy. *J Pain Symptom Manage* 29: 401-410.

9. Goodman LS, Limbird LE, Milinoff PB (1996) (ed) Goodman and Gilman's The pharmacological basis of therapeutics. 9th ed New York NY: McGraw-Hill Professional.
10. Rao RD, Michalak JC, Sloan JA, Loprinzi CL, Soori GS, et al. (2007) Efficacy of gabapentin in the management of chemotherapy-induced peripheral neuropathy: a phase 3 randomized, double-blind, placebo-controlled, crossover trial (N00C3).. *Cancer* 110: 2110-2118.
11. Suzuki T, Nishio K, Tanabe S (2001) The MRP family and anticancer drug metabolism. *Curr Drug Metab* 2: 367-377.
12. Gustafsson T, Puntschart A, Kaijser L, Jansson E, Sundberg CJ (1999) Exercise-induced expression of angiogenesis-related transcription and growth factors in human skeletal muscle. *Am J Physiol* 276: H679-H685.
13. Fukai T, Siegfried MR, Ushio-Fukai M, Cheng Y, Kojda G, et al. (2000) Regulation of the vascular extracellular superoxide dismutase by nitric oxide and exercise training. *J Clin Invest* 105: 1631-1639.
14. Markes M, Brockow T, Resch KL (2006) Exercise for women receiving adjuvant therapy for breast cancer. *Cochrane Database Syst Rev* CD005001.
15. Irwin ML, Crumley D, McTiernan A, Bernstein L, Baumgartner R, et al. (2003) Physical activity levels before and after a diagnosis of breast carcinoma: the Health, Eating, Activity, and Lifestyle (HEAL) study. *Cancer* 97: 1746-1757.
16. Herrero F, San Juan AF, Fleck SJ, Balmer J, Perez M, et al. (2006) Combined aerobic and resistance training in breast cancer survivors: A randomized, controlled pilot trial. *Int J Sports Med* 27: 573-580.
17. Aitken DR, James AG (1997) Seromas and physiotherapy after mastectomy. *Ann Surg Oncol* 4: 293-297.
18. Schneider CM, Dennehy CA, Roozeboom M, Carter SD (2002) A model program: exercise intervention for cancer rehabilitation. *Integr Cancer Ther* 1: 76-82.
19. Schneider CM, Hsieh CC, Sprod LK, Carter SD, Hayward R (2007) Effects of supervised exercise training on cardiopulmonary function and fatigue in breast cancer survivors during and after treatment. *Cancer* 110: 918-925.
20. Schneider CM, Hsieh CC, Sprod LK, Carter SD, Hayward R (2007) Exercise training manages cardiopulmonary function and fatigue during and following cancer treatment in male cancer survivors. *Integr Cancer Ther* 6: 235-241.
21. Schneider CM, Hsieh CC, Sprod LK, Carter SD, Hayward R (2007) Cancer treatment-induced alterations in muscular fitness and quality of life: the role of exercise training. *Ann Oncol* 18: 1957-1962.
22. Balducci S, Iacobellis G, Parisi L, Di Biase N, Calandriello E, et al. (2006) Exercise training can modify the natural history of diabetic peripheral neuropathy. *J Diabetes Complications* 20: 216-223.
23. Richardson JK, Sandman D, Vela S (2001) A focused exercise regimen improves clinical measures of balance in patients with peripheral neuropathy. *Arch Phys Med Rehabil* 82: 205-209.
24. Tesfaye S, Harris ND, Wilson RM, Ward JD (1992) Exercise-induced conduction velocity increment: A marker of impaired peripheral nerve blood flow in diabetic neuropathy. *Diabetologia* 35: 155-159.
25. Cohen SR, Mount BM, Strobel MG, Bui F (1995) The McGill Quality of Life Questionnaire: a measure of quality of life appropriate for people with advanced disease. A preliminary study of validity and acceptability. *Palliat Med* 9: 207-219.
26. Bennett M (2001) The LANSS Pain Scale: the Leeds assessment of neuropathic symptoms and signs. *Pain* 92: 147-157.
27. Centers for Disease Control and Prevention (2011) How much physical activity do adults need?
28. del Porto LA, Nicholson GA, Ketheswaren P (2010) Correlation between muscle atrophy on MRI and manual strength testing in hereditary neuropathies. *J Clin Neurosci* 17: 874-878.
29. Carter GT, Abresch RT, Fowler WM Jr, Johnson ER, Kilmer DD, et al. (1995) Profiles of neuromuscular diseases. Hereditary motor and sensory neuropathy, types I and II. *Am J Phys Med Rehabil* 74: S140-S149.
30. Lindeman E, Leffers P, Spaans F, Drukker J, Reulen J, et al. (1995) Strength training in patients with myotonic dystrophy and hereditary motor and sensory neuropathy: a randomized clinical trial. *Arch Phys Med Rehabil* 76: 612-620.
31. Kilmer DD, McCrory MA, Wright NC, Aitkens SG, Bernauer EM, et al. (1994) The effect of a high resistance exercise program in slowly progressive neuromuscular disease. *Arch Phys Med Rehabil* 75: 560-563.
32. Fisher MA, Langbein WE, Collins EG, Williams K, Corzine L (2007) Physiological improvement with moderate exercise in type II diabetic neuropathy. *Electromyogr Clin Neurophysiol* 47: 23-28.
33. Smith AG, Russell J, Feldman EL, Goldstein J, Peltier A, et al. (2006) Lifestyle intervention for pre-diabetic neuropathy. *Diabetes Care* 29: 1294-1299.
34. Spellman CW (2008) Aggressively managing type 2 diabetes mellitus, hyperlipidemia, and bone loss. *J Am Osteopath Assoc* 108: S20-S27.
35. Van meeteren NL, Brakkee JH, Biessels GJ, Kappelle AC, Helders PJ et al. (1996) Effect of exercise training on acute (crush lesion) and chronic (diabetes mellitus) peripheral neuropathy in the rat. *Restor Neurol Neurosci* 10: 85-93.
36. Cook DB, Nagelkirk PR, Poluri A, Mores J, Natelson BH (2006) The influence of aerobic fitness and fibromyalgia on cardiorespiratory and perceptual responses to exercise in patients with chronic fatigue syndrome. *Arthritis Rheum* 54: 3351-3362.
37. Cook DB, Stegner AJ, Ellingson LD (2010) Exercise alters pain sensitivity in Gulf War Veterans with chronic musculoskeletal pain. *J Pain* 11: 764-772.
38. Schenone A, Nobbio L, Monti Bragadin M, Ursino G, Grandis M (2011) Inherited neuropathies. *Curr Treat Options Neurol* 13: 160-179.
39. Udina E, Puigdemasa A, Navarro X (2011) Passive and active exercise improve regeneration and muscle reinnervation after peripheral nerve injury in the rat. *Muscle Nerve* 43: 500-509.
40. Seo TB, Oh MJ, You BG, Kwon KB, Chang IA, et al. (2009) ERK 1/2-mediated Schwann cell proliferation in the regenerating sciatic nerve by treadmill training. *J Neurotrauma* 26: 1733-1744.
41. Portenoy RK, Hagen NA (1990) Breakthrough pain: definition, prevalence and characteristics. *Pain* 41: 273-281.
42. Koltyn KF, Garvin AW, Gardiner RL, Nelson TF (1996) Perception of pain following aerobic exercise. *Med Sci Sports Exerc* 28: 1418-1421.
43. Gurevich M, Kohn PM, Davis C (1994) Exercise-induced analgesia and the role of reactivity in pain sensitivity. *J Sports Sci* 12: 549-559.
44. Rief W, Bardwell WA, Dimsdale JE, Natarajan L, Flatt SW, et al. (2011) Long-term course of pain in breast cancer survivors: a 4-year longitudinal study. *Breast Cancer Res Treat* 130: 579-586.
45. Liou YM, Lee HL, Chien LY, Kao WY, Chiang CC, et al. (2011) Daily-life physical activity and related factors among patients with cancer receiving chemotherapy in Taiwan. *Cancer Nurs* 34: 443-452.
46. Courneya KS, Tamburrini AL, Woolcott CG, McNeely ML, Karvinen KH, et al. (2011) The Alberta physical activity and breast cancer prevention trial: Quality of life outcomes. *Prev Med* 52: 26-32.
47. Griffith K, Wenzel J, Shang J, Thompson C, Stewart K, et al. (2009) Impact of a walking intervention on cardiorespiratory fitness, self-reported physical function, and pain in patients undergoing treatment for solid tumors. *Cancer* 115: 4874-4884.
48. Fink A (1995) How to sample in surveys. Thousand Oaks, CA: SAGE.
49. Fowler FJ (1993) Survey research methods (2nd ed.). Thousand Oaks, CA: SAGE.

Submit your next manuscript and get advantages of OMICS Group submissions

Unique features:

- User friendly/feasible website-translation of your paper to 50 world's leading languages
- Audio Version of published paper
- Digital articles to share and explore

Special features:

- 200 Open Access Journals
- 15,000 editorial team
- 21 days rapid review process
- Quality and quick editorial, review and publication processing
- Indexing at PubMed (partial), Scopus, DOAJ, EBSCO, Index Copernicus and Google Scholar etc
- Sharing Option: Social Networking Enabled
- Authors, Reviewers and Editors rewarded with online Scientific Credits
- Better discount for your subsequent articles

Submit your manuscript at: <http://www.editorialmanager.com/acrgroup/>