

7-2009

# The Effects of an Acute Dose of *Rhodiola rosea* on Exercise Performance and Cognitive Function

Eric E. Noreen  
*Gettysburg College*

James G. Buckley  
*Gettysburg College*

Stephanie L. Lewis  
*Gettysburg College*

## **Roles**

James G. Buckley: Class of 2009  
Stephanie L. Lewis: Class of 2009

Follow this and additional works at: <https://cupola.gettysburg.edu/healthfac>

Part of the [Other Medicine and Health Sciences Commons](#), and the [Sports Sciences Commons](#)

**Share feedback about the accessibility of this item.**

---

Noreen, Eric, James G. Buckley, and Stephanie L. Lewis. "The Effects of an Acute Dose of *Rhodiola rosea* on Exercise Performance and Cognitive Function." *Journal of the International Society of Sports Nutrition* 6 Supplement I (July 2009), 14.

This is the publisher's version of the work. This publication appears in Gettysburg College's institutional repository by permission of the copyright owner for personal use, not for redistribution. Cupola permanent link: <https://cupola.gettysburg.edu/healthfac/53>

This open access article is brought to you by The Cupola: Scholarship at Gettysburg College. It has been accepted for inclusion by an authorized administrator of The Cupola. For more information, please contact [cupola@gettysburg.edu](mailto:cupola@gettysburg.edu).

---

# The Effects of an Acute Dose of *Rhodiola rosea* on Exercise Performance and Cognitive Function

## Abstract

**Background:** The purpose of this study was to determine the effects of an acute oral dose of 3 mg/kg of *Rhodiola rosea* (*R. rosea*) on endurance exercise performance, mood, and cognitive function.

**Methods:** A total of 15 recreationally active college women ( $21.3 \pm 0.09$  y,  $56.1 \pm 6.3$  kg; mean  $\pm$  SD) participated in this study. 2–7 d after a familiarization trial subjects ingested in a double blind, random crossover manner, either *R. rosea* or a carbohydrate placebo 1 h prior to testing. Exercise testing consisted of a 10 minute warm-up, standardized to 80% of the average watts produced during the familiarization trial, followed by a 6 mile simulated indoor time trial on a Velotron electronic bicycle ergometer. Every 5 min during the time trial, subjects rated their level of perceived exertion using a BORG 10 pt scale. A blood sample was taken pre warm-up, 2 minutes post warm-up, and 2 minutes following completion of the time trial, and was analyzed for lactate concentration. Subjects also completed a Profile of Mood States (POMS) questionnaire and a Stroop's color test pre-warm up and following the completion of the time trial. Subjects returned to the lab 2–7 d later to repeat the testing with the other condition.

**Results:** A 3 mg/kg acute dose of *R. rosea* resulted in a shorter time to completion of the 6 mile time trial course (*R. rosea*  $1544.7 \pm 155.2$  s, Placebo  $1569.5 \pm 179.4$  s; mean  $\pm$  SD;  $p = 0.06$ ) as well as a lower average heart rate during the standardized warm up (*R. rosea*  $138.6 \pm 13.3$  bpm, Placebo  $143.7 \pm 12.4$  bpm; mean  $\pm$  SD;  $p = 0.001$ ). There were no significant differences between treatment conditions for rating of perceived exertion during the time trial. Both treatments resulted in a significant increase in the POMS fatigue score following exercise ( $p = 0.001$ ), as well as a significant improvement following exercise for the Stroop's test of incongruent words ( $p = 0.001$ ). No other significant differences between treatments were observed.

**Conclusion:** Acute *Rhodiola rosea* ingestion decreases the heart rate response to sub-maximal exercise, and appears to improve endurance exercise performance.

## Keywords

Sports Science, *Rhodiola rosea*, Herbal Supplement

## Disciplines

Other Medicine and Health Sciences | Sports Sciences

## Comments

This article is part of the supplement: [Proceedings of the Sixth International Society of Sports Nutrition \(ISSN\) Conference and Expo](#)

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Poster presentation

## The effects of an acute dose of *Rhodiola rosea* on exercise performance and cognitive function

Eric Noreen\*, James Buckley and Stephanie Lewis

Address: The Department of Health Sciences, Gettysburg College, Gettysburg, PA 17325, USA

Email: Eric Noreen\* - enoreen@gettysburg.edu

\* Corresponding author

from 2009 International Society of Sports Nutrition Conference and Expo  
New Orleans, LA, USA. 14–15 June 2009

Published: 31 July 2009

Journal of the International Society of Sports Nutrition 2009, 6(Suppl 1):P14 doi:10.1186/1550-2783-6-S1-P14

This abstract is available from: <http://www.jissn.com/content/6/S1/P14>

© 2009 Noreen et al; licensee BioMed Central Ltd.

### Background

The purpose of this study was to determine the effects of an acute oral dose of 3 mg/kg of *Rhodiola rosea* (*R. rosea*) on endurance exercise performance, mood, and cognitive function.

### Methods

A total of 15 recreationally active college women ( $21.3 \pm 0.09$  y,  $56.1 \pm 6.3$  kg; mean  $\pm$  SD) participated in this study. 2–7 d after a familiarization trial subjects ingested in a double blind, random crossover manner, either *R. rosea* or a carbohydrate placebo 1 h prior to testing. Exercise testing consisted of a 10 minute warm-up, standardized to 80% of the average watts produced during the familiarization trial, followed by a 6 mile simulated indoor time trial on a Velotron electronic bicycle ergometer. Every 5 min during the time trial, subjects rated their level of perceived exertion using a BORG 10 pt scale. A blood sample was taken pre warm-up, 2 minutes post warm-up, and 2 minutes following completion of the time trial, and was analyzed for lactate concentration. Subjects also completed a Profile of Mood States (POMS) questionnaire and a Stroop's color test pre-warm up and following the completion of the time trial. Subjects returned to the lab 2–7 d later to repeat the testing with the other condition.

### Results

A 3 mg/kg acute dose of *R. rosea* resulted in a shorter time to completion of the 6 mile time trial course (*R. rosea*

$1544.7 \pm 155.2$  s, Placebo  $1569.5 \pm 179.4$  s; mean  $\pm$  SD;  $p = 0.06$ ) as well as a lower average heart rate during the standardized warm up (*R. rosea*  $138.6 \pm 13.3$  bpm, Placebo  $143.7 \pm 12.4$  bpm; mean  $\pm$  SD;  $p = 0.001$ ). There were no significant differences between treatment conditions for rating of perceived exertion during the time trial. Both treatments resulted in a significant increase in the POMS fatigue score following exercise ( $p = 0.001$ ), as well as a significant improvement following exercise for the Stroop's test of incongruent words ( $p = 0.001$ ). No other significant differences between treatments were observed.

### Conclusion

Acute *Rhodiola rosea* ingestion decreases the heart rate response to sub-maximal exercise, and appears to improve endurance exercise performance.