



Fall 2015

Isolation & Characterization of Bacteria in the Built Environment: Measuring The Effect of Pharmaceuticals on Growth

Amanda V. Finck
Gettysburg College

Kelly J. DiGeronimo
Gettysburg College

Nene S. Sy
Gettysburg College

Zakiya Whatley
Gettysburg College

Follow this and additional works at: https://cupola.gettysburg.edu/student_scholarship

 Part of the [Bacteriology Commons](#), and the [Biology Commons](#)

Share feedback about the accessibility of this item.

Finck, Amanda V.; DiGeronimo, Kelly J.; Sy, Nene S.; and Whatley, Zakiya, "Isolation & Characterization of Bacteria in the Built Environment: Measuring The Effect of Pharmaceuticals on Growth" (2015). *Student Publications*. 382.
https://cupola.gettysburg.edu/student_scholarship/382

This is the author's version of the work. This publication appears in Gettysburg College's institutional repository by permission of the copyright owner for personal use, not for redistribution. Cupola permanent link: https://cupola.gettysburg.edu/student_scholarship/382

This open access conference material is brought to you by The Cupola: Scholarship at Gettysburg College. It has been accepted for inclusion by an authorized administrator of The Cupola. For more information, please contact cupola@gettysburg.edu.

Isolation & Characterization of Bacteria in the Built Environment: Measuring The Effect of Pharmaceuticals on Growth

Abstract

This work reports the isolation and characterization of bacteria from the built environment at Gettysburg College in Gettysburg, PA. Surfaces of a water fountain on campus were swabbed and serially streaked to isolate multiple bacteria on R2A agar. Following multiple rounds of growth, the unknown microbial candidates were narrowed to two visibly distinct organisms. Morphological characterization and phylogenetic identification based on 16S rDNA sequencing revealed that the isolates were *Chryseobacterium hispalense* and *Microbacterium maritopicum*. We report synergistic biofilm formation between *Chryseobacterium hispalense* and *Microbacterium maritopicum*. The contamination of drinking water with varying levels of personal care products and pharmaceuticals (PCPPs) is well documented. Additionally, these environmental pollutants and their derivatives affect aquatic life, as illustrated with effect of the antidepressant fluoxetine on mudsnails. To determine if previously reported contaminants affect freshwater bacteria, we assessed both planktonic growth and biofilm formation following exposure to nalidixic acid (nonfluorinated quinolone antibiotic), diphenhydramine (over-the-counter drug Benadryl), and fluoxetine (Prozac).

Keywords

Chryseobacterium hispalense, *Microbacterium maritopicum*, personal care products and pharmaceuticals, fluoxetine, nalidixic acid, synergistic biofilm formation, biofilm formation, Microbiology

Disciplines

Bacteriology | Biology | Microbiology

Comments

HHMI Grant Funding

This poster was presented at the [18th Undergraduate Research Symposium in the Chemical and Biological Sciences at UMBC](#), October 2015.



Isolation & Characterization of Bacteria in the Built Environment: Measuring the Effect of Personal Care Products & Pharmaceuticals on Growth



Amanda Finck¹, Kelly DiGeronimo¹, Nene Sy¹, Zakiya Whatley¹

¹ Department of Biology, Gettysburg College, 300 N Washington Street, Gettysburg, PA 17325
Funding Provided by HHMI Grant #52007540 – Cross Disciplinary Sciences at Gettysburg College

BACKGROUND

The contamination of drinking water with varying levels of personal care products and pharmaceuticals (PCPPs) is well documented. Additionally, these environmental pollutants and their derivatives affect aquatic life, as illustrated with effect of the antidepressant fluoxetine on molluscs and crustaceans (Fong and Ford 2014). To determine if previously reported contaminants affect freshwater bacteria, we isolated and characterized unknown bacteria from the built environment at Gettysburg College in Gettysburg, PA. Additionally, we measured both planktonic growth and biofilm formation.

ISOLATION OF MICROBES




Figure 1. A) Microbes were isolated from the drain of a water fountain in an academic building (McCreary Hall) at Gettysburg College. B) Swabs from sample were streaked and grown on R2A agar at room temperature for 3 days.

MOLECULAR IDENTIFICATION




Figure 2. Unknown microbes were identified via PCR amplification of 16S rDNA and subsequent sequencing. Colony PCR with primers U1 (5'-ACGCGTCGACAGAGTTGATCTGGCT-3') and U2 (5'-CGCGGATCCGCTACCTTGTACGACT-3') yielded ~1500 bp PCR products. PCR products were visualized on a 1% agarose gel, shown left. See amplification of desired 16S rDNA isolated species (Wells 2-4) and absence of product in the Negative Control (Well 5).




Figure 3: PCR products shown in Figure 2 were sequenced by Eton Biolabs using primers U1 and U2. Representative results are shown above.

Genus Species	WFW	e value	Percent Identity	Length (bp)
<i>Microbacterium maritopicum</i> strain DSM 12512	ref NR_114986.1	0.0	910/910(100%)	1345
<i>Microbacterium maritopicum</i> strain DSM 12512	ref NR_042351.1	0.0	910/910(100%)	1437
<i>Microbacterium oxydans</i> strain DSM 20578	ref NR_044931.1	0.0	909/910(99%)	1466
Genus Species	WFO	e value	Percent Identity	Length (bp)
<i>Chryseobacterium hispalense</i> strain AG13	ref NR_116277.1	0.0	769/769(100%)	1480
<i>Chryseobacterium taeanaense</i> strain NBRC 100863	ref NR_113951.1	0.0	745/771(97%)	1442
<i>Chryseobacterium taeanaense</i> strain PHA3-4	ref NR_043254.1	0.0	745/771(97%)	1424

Table 1. Phylogenetic Analysis of 16S rDNA. Using NCBI Blast, sequences were compared to a database of 16S rDNA to determine the closest matches.

RESEARCH QUESTION

We ask what bacteria are present in the built environment? Additionally, how are these organisms affected by pharmaceutical and personal care products known to contaminate drinking water?




Figure 5. *M. maritopicum* and *C. hispalense* grow @ 37°C. Overnight cultures were diluted 100 fold in R2A broth and incubated with shaking every 780 seconds. Growth was measured in a 96-well microtiter plate, with OD595 recorded every 15 minutes for 18 hours.




Figure 6. At high concentrations ($5 \times 10^{-3} M - 5 \times 10^{-4} M$), fluoxetine completely inhibits growth. Growth defects are also seen at more dilute concentrations ($5 \times 10^{-5} - 5 \times 10^{-6} M$). Overnight cultures were diluted and growth was monitored as explained above. Results are from 3 separate trials, n=14.




Figure 7. Nalidixic acid impairs growth at higher concentrations (4 µg/mL - 1.44 µg/mL).

Antibiotic	<i>Chryseobacterium hispalense</i>	<i>Microbacterium maritopicum</i>
Rifampin	Susceptible 28 mm	Resistant 14 mm
Erythromycin	Susceptible 21 mm	Susceptible 19 mm
Vancomycin	Susceptible 20 mm	Susceptible 25 mm
Tetracycline	Susceptible 24 mm	Resistant 8 mm
Penicillin	Susceptible 15 mm	Susceptible 27 mm

Table 2. Antibiotic resistance or susceptibility was determined using a standard filter disk assay on Mueller-Hinton agar. Following overnight incubation, zones of inhibition were measured and categorized based on Clinical Laboratory and Standards Institutes regulations.

EXPERIMENTAL DESIGN





Figure 8. *C. hispalense* and *M. maritopicum* exhibit synergistic biofilm formation. This synergism appears specific to these species, as *C. hispalense* & *E. coli* co-incubation does not yield biofilm at similar levels. Static cultures were incubated 3 days @ room temperature. Planktonic bacteria was removed and biofilm was stained with crystal violet. Crystal violet was solubilized in 30% acetic acid and measured at 595nm. Results represent 1 trial, n=5.

CONCLUSIONS

We identified two unknown microbes as *Microbacterium maritopicum* and *Chryseobacterium hispalense*. Identification was confirmed by phenotypic & phylogenetic analysis of 16S rDNA (Fig 2-3 & Table 1). We found *M. maritopicum* was resistant to rifampin and tetracycline, while *C. hispalense* was susceptible to all antibiotics tested (Table 2). We report a previously uncharacterized growth impairment in response to fluoxetine, a serotonin reuptake inhibitor (Figure 6). This finding was interesting and unexpected. Additionally, abiotic biofilm assays indicate specific synergism between *C. hispalense* and *M. maritopicum* (Figure 8).

Future experiments include additional tests to understand the effect of fluoxetine and other known PCPP contaminants on freshwater bacteria. We are particularly interested in the effects of these compounds on biofilm formation. Does drug exposure shift the composition of biofilms? How does drug exposure affect gene transfer within these communities? Do trace levels of combinations of drugs create a measurable response? These questions are particularly important when considering how contamination affects both drinking water and freshwater ecosystems.

REFERENCES

Fong P, Ford A. The biological effects of antidepressants on the molluscs and crustaceans: a review. *Aquat Toxicol.* 2014;151:4-13.